



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MAY 7.

ARTHUR ERWIN BROWN, D.Sc., Vice-President, in the Chair.

Twenty-one persons present.

The Chair announced the death of William K. Shryock, a member, April 28, 1907.

The Publication Committee reported that papers under the following titles had been presented for publication:

"Description of New Species of Spioniform Annelids," by J. Percy Moore (April 26).

"A Contribution to the Morphology of *Pyrula*," by Burnett Smith (April 29).

"The Hymenopterous Family *Evaniidæ*," by J. Chester Bradley (April 3).

Origin of the Tropical Forms of the Land Molluscan Fauna of Southern Florida.—DR. H. A. PILSBRY remarked that former writers on this fauna have treated the tropical area in Florida as an outlier of the Antillean zoological province. The speaker recognized two groups of genera, of diverse origin: (1) Antillean genera, consisting chiefly of species also West Indian, such as *Chondropoma*, *Liguus*, *Cepolis*, *Varicella*, etc., concentrated southward, and unknown in the Floridian Pliocene, and (2) Mexican genera, chiefly composed of species confined to Florida, such as *Euglandina*, *Praticolella*, *Drymæus* of the *dormani* type, and apparently *Polygyra*, some of which are represented in the Floridian (Caloosahatchie) Pliocene. The first group he regarded as recent additions to the fauna, which had reached Florida from Cuba and the Bahamas by the agency of hurricanes, drifting trees and the like. The second group probably emigrated from the Southwest during the mild Miocene climate, and entered peninsular Florida on the elevation marking the close of the Miocene. The Oligocene fauna of the Tampa Siliceous beds, the speaker thought, had left no descendants. Its snails were of wholly Antillean type, the leading Antillean genera of *Helices*, *Cepolis* and *Pleurodonte* both being represented.

Dr. Pilsbry alluded also to Prof. Alexander Agassiz's conclusion, that the Florida Keys are in process of diminution by solution of the limestone composing them. The former unity of keys now separated accounts for the homogeneity of their land-shell faunas.